

$B_{s,d} \rightarrow \mu^+ \mu^-$ in technicolor model with scalars

Zhaohua Xiong^{1,2} and Jin Min Yang³

¹ *CCAST (World Laboratory), P. O. Box 8730, Beijing 100080, China*

² *Institute of High Energy Physics, Academia Sinica, Beijing 100039, China*

³ *Institute of Theoretical Physics, Academia Sinica, Beijing 100080, China*

(Dated: February 1, 2008)

Rare decays $B_{s,d} \rightarrow \mu^+ \mu^-$ are evaluated in technicolor model with scalars. R_b is revisited to constrain the model parameter space. It is found that restriction on f/f' arising from R_b which was not considered in previous studies requires f/f' no larger than 1.9 at 95% confidence level, implying no significantly enhancement for $Br(B_{s,d} \rightarrow \mu^+ \mu^-)$ from neutral scalars in the model. However, the branching ratio of $B_s \rightarrow \mu^+ \mu^-$ can still be enhanced by a factor of 5 relative to the standard model prediction. With the value of $f/f' \lesssim 1.9$, an upgraded Tevatron with an integrated luminosity $20fb^{-1}$ will be sensitive to enhancement of $B_s \rightarrow \mu^+ \mu^-$ in this model provided that neutral scalar mass m_σ is below 580 GeV.

PACS numbers: 12.60.Nz, 13.20.Hw, 13.38.Dg

I. INTRODUCTION

The flavor-changing neutral-current B-meson rare decays play an important role for testing the Standard Model (SM) at loop level and probing new physics beyond the SM. Among these decays, $B_{s,d} \rightarrow \mu^+ \mu^-$ are of special interest due to their relative cleanliness and good sensitivity to new physics.

There are numerous speculations on the possible forms of new physics, among which supersymmetry and technicolor are the two typical different frameworks. Both frameworks are well motivated. As a low-energy effective theory, the technicolor model with scalars introduces additional scalars to connect the technicolor condensate to the ordinary fermions [1]. The phenomenology of this model has been considered extensively in the literature [1, 2, 3, 4, 5, 6, 7, 8]. It has been found that this model does not produce unacceptably large contributions to neutral meson mixings or to the electroweak S and T parameters [1, 2]. On the other hand, this model does predict potentially visible contributions to b-physics observables such as R_b [6] and the rate of various rare B-meson decays [6, 7, 8].

Studies [9] showed that the processes $B_{s,d} \rightarrow \mu^+ \mu^-$ are sensitive to supersymmetry. In this Letter we will extend our previous studies [8, 10] and evaluate the branching ratio of $B_{s,d} \rightarrow \mu^+ \mu^-$ in the technicolor model with scalars. First we will present a brief description of the model, then give the analytical calculations for $B_{s,d} \rightarrow \mu^+ \mu^-$. We will focus our attention on the neutral scalars contributions, which are likely to be sizable because, as shown in our following analysis, they will be enhanced by a factor $(f/f')^4$ as the parameter f/f' gets large. Before performing the numerical calculations, we examine the current bounds on this model from a variety of experiments, especially the latest measurements of R_b [11]. Since the theoretical expression for R_b used in constraining the model parameter space [6] seems not right, we will recalculate the contributions to R_b from the scalars in this model. We find the constraint from R_b is still strongest

as indicated in [20], compared with those from the direct searches for neutral and charged scalars [12], $B^0 - \bar{B}^0$ mixing, $b \rightarrow s\gamma$ [13] as well as the muon anomalous magnetic moment [14]. Further, we evaluate restriction on f/f' arising from R_b which was not considered in previous studies. Subject to the current bounds, the numerical results are presented in Sec. V. Finally, the conclusion is assigned in Sec. VI.

II. THE TECHNICOLOR MODEL WITH SCALARS

In this section we will briefly discuss the technicolor model with scalars and give the relevant Lagrangians which are needed in our calculations. More details of the model have been described in Refs. [1, 2].

The model embraces the full SM gauge structure and all SM fermions which are technicolor singlets. It has a minimal $SU(N)$ technicolor sector, with two techniflavors that transform as a left-handed doublet and two right-handed singlets under $SU(2)_W$,

$$T_L = \begin{pmatrix} p \\ m \end{pmatrix}_L, \quad p_R, m_R \quad (1)$$

with weak hypercharges $Y(T_L) = 0$, $Y(p_R) = 1$, and $Y(m_R) = -1$. All of the fermions couple to a weak scalar doublet ϕ to which both the ordinary fermions and technifermions are coupled. This scalar's purpose is to couple the technifermion condensate to the ordinary fermions and thereby generate fermion masses. If we write the matrix form of the scalar doublet as

$$\Phi = \begin{bmatrix} \bar{\phi}^0 & \phi^+ \\ -\phi^- & \phi^0 \end{bmatrix} \equiv \frac{(\sigma + f')}{\sqrt{2}} \Sigma', \quad (2)$$

and adopt the non-linear representation $\Sigma = \exp(\frac{2i\Pi}{f})$ and $\Sigma' = \exp(\frac{2i\Pi'}{f'})$ for technipion, with fields in Π and Π' representing the pseudoscalar bound states of the technifermions p and m , then the kinetic terms for the scalar

fields are given by

$$\begin{aligned} \mathcal{L}_{K.E.} = & \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma + \frac{1}{4} f^2 \text{Tr}(D_\mu \Sigma^\dagger D^\mu \Sigma) \\ & + \frac{1}{4} (\sigma + f')^2 \text{Tr}(D_\mu \Sigma'^\dagger D^\mu \Sigma'). \end{aligned} \quad (3)$$

Here D^μ (D'^μ) denote the $SU(2)_L \times SU(2)_R$ covariant derivatives, σ is an isosinglet scalar field, f and f' are the technipion decay constant and the effective vacuum expectation value (VEV), respectively.

As mixing between Π and Π' occurs, π_a and π_p are formed with π_a becoming the longitudinal component of the W and Z, and π_p remaining in the low-energy theory as an isotriplet of physical scalars. From Eq. (3) one can obtain the correct gauge boson masses providing that $f^2 + f'^2 = v^2$ with the electroweak scale $v = 246 \text{ GeV}$.

Additionally, the contributions to scalar potential generated by the technicolor interactions should be included in this model. The simplest term one can construct is

$$\mathcal{L}_T = c_1 4\pi f^3 \text{Tr} \left[\Phi \begin{pmatrix} h_+ & 0 \\ 0 & h_- \end{pmatrix} \Sigma^\dagger \right] + h.c., \quad (4)$$

where c_1 is a coefficient of order unity, h_+ and h_- are the Yukawa couplings of scalars to p and m . From Eq. (4) the mass of the charged scalar at lowest order is obtained as

$$m_{\pi_p}^2 = 2c_1 \sqrt{2} \frac{4\pi f}{f'} v^2 h \quad (5)$$

with $h \equiv (h_+ + h_-)/2$. To absorb the largest Coleman-Weinberg radiative corrections [15] for the σ field which affect the phenomenology of the charged scalar, the shifted scalar mass $\widetilde{M}_\phi$ and coupling $\tilde{\lambda}$ are determined by

$$\widetilde{M}_\phi^2 f' + \frac{\tilde{\lambda}}{2} f'^3 = 8\sqrt{2} c_1 \pi h f^3. \quad (6)$$

Therefore, the mass of the scalar σ can be expressed as

$$m_\sigma^2 = \widetilde{M}_\phi^2 + \frac{2}{3\pi^2} \left[6 \left(\frac{m_t}{f'} \right)^4 + N h^4 \right] f'^2 \quad (7)$$

in limit (i) where the shifted ϕ^4 coupling $\tilde{\lambda}$ is small and can be neglected and

$$m_\sigma^2 = \frac{3}{2} \tilde{\lambda} f'^2 - \frac{1}{4\pi^2} \left[6 \left(\frac{m_t}{f'} \right)^4 + N h^4 \right] f'^2. \quad (8)$$

in limit (ii) where the shifted mass of the scalar doublet ϕ , $\widetilde{M}_\phi$ is small and can be neglected. The advantage of this model is at the lowest order, only two independent parameters in the limits (i) and (ii) are needed to describe the phenomenology. We choose (h, m_σ) as physical parameters and assume $N = 4$ and $c_1 = 1$ in numerical calculations.

III. CALCULATIONS

We start the calculation by writing down the effective Hamiltonian describing the process $B_q \rightarrow \mu^+ \mu^-$ ($q=s,d$)

$$\begin{aligned} \mathcal{M} = & \frac{\alpha G_F}{\sqrt{2}\pi} V_{tb} V_{tq}^* \left\{ -2C_7^{eff} \frac{m_b}{p^2} \bar{q} i \sigma_{\mu\nu} p^\nu P_R b \right. \\ & + C_9^{eff} \bar{q} \gamma_\mu P_L b \bar{\mu} \gamma^\mu \mu + C_{10} \bar{q} \gamma_\mu P_L b \bar{\mu} \gamma^\mu \gamma_5 \mu \\ & \left. + C_{Q_1} \bar{q} P_R b \bar{\mu} \mu + C_{Q_2} \bar{q} P_R b \bar{\mu} \gamma_5 \mu \right\}, \end{aligned} \quad (9)$$

where $P_{R,L} = \frac{1}{2}(1 \pm \gamma_5)$, p is the momentum transfer. Operators $\mathcal{O}_{7,9,10}$ which correspond to the first three Wilson coefficients are the same as those given in [16] and $\mathcal{Q}_{1,2}$ corresponding to the last two are the additional operators arising from the neutral scalars exchange diagrams [17].

Using the effective Hamiltonian and

$$\begin{aligned} \langle 0 | \bar{q} \gamma_\mu \gamma_5 b | B_q \rangle &= -f_{B_q} p_\mu, \\ \langle 0 | \bar{q} \gamma_5 b | B_q \rangle &= -f_{B_q} m_{B_q}, \\ \langle 0 | \bar{q} \sigma_{\mu\nu} (1 + \gamma_5) b | B_q \rangle &= 0, \end{aligned} \quad (10)$$

we find that only operator $\mathcal{O}_{10}$ and $\mathcal{Q}_{1,2}$ contribute to process $B_q \rightarrow \mu^+ \mu^-$ with the decay rate given by

$$\begin{aligned} \Gamma(B_q \rightarrow \mu^+ \mu^-) &= \frac{\alpha^2 G_F^2}{64\pi^3} |V_{tb} V_{tq}^*|^2 f_{B_q}^2 m_{B_q}^3 \\ &\times \left[C_{Q_1}^2 + \left(C_{Q_2} + \frac{2m_\mu}{m_{B_q}} C_{10} \right)^2 \right]. \end{aligned} \quad (11)$$

For convenience, we write down the branching fractions numerically

$$\begin{aligned} Br(B_d \rightarrow \mu^+ \mu^-) &= 3.8 \times 10^{-9} \left[\frac{\tau_{B_d}}{1.65 \text{ ps}} \right] \left[\frac{f_{B_d}}{210 \text{ MeV}} \right]^2 \\ &\times \left| \frac{V_{td}}{0.008} \right|^2 \left[\frac{m_{B_d}}{5.28 \text{ GeV}} \right]^3 \left[C_{Q_1}^2 \right. \\ &\left. + \left(C_{Q_2} + 2 \frac{m_\mu}{m_{B_d}} C_{10} \right)^2 \right], \\ Br(B_s \rightarrow \mu^+ \mu^-) &= 1.2 \times 10^{-7} \left[\frac{\tau_{B_s}}{1.49 \text{ ps}} \right] \left[\frac{f_{B_s}}{245 \text{ MeV}} \right]^2 \\ &\times \left| \frac{V_{ts}}{0.04} \right|^2 \left[\frac{m_{B_s}}{5.37 \text{ GeV}} \right]^3 \left[C_{Q_1}^2 \right. \\ &\left. + \left(C_{Q_2} + 2 \frac{m_\mu}{m_{B_s}} C_{10} \right)^2 \right], \end{aligned} \quad (12)$$

where τ_{B_q} and f_{B_q} are the B_q lifetime and decay constant, respectively.

In the technicolor model with scalars, the additional contributions arise from the scalars. The contributions of the charged scalar $\pi_p^\pm$ with gauge boson Z, γ exchanges to the Wilson coefficients C_{10} at m_W scale have been calculated by using Feynman rules derived from Eq. (3), (4)

and given by [7, 8]

$$C_{10}(m_W)_{TC} = \frac{x_W}{\sin^2 \theta_W} \left(\frac{f}{f'} \right)^2 \left[-\frac{x_{\pi_p}}{8(x_{\pi_p} - 1)} + \frac{x_{\pi_p}}{8(x_{\pi_p} - 1)^2} \ln x_{\pi_p} \right], \quad (13)$$

where θ_W is the Weinberg angle and $x_i = m_i^2/m_t^2$. As for the contributions arising from the neutral scalars exchanges, when only the leading terms in large f/f' limit kept, they can be expressed as [8]

$$\begin{aligned} C_{Q_1}(m_W)_{TC} &= -\frac{x_W}{\sin^2 \theta_W} \left(\frac{f}{f'} \right)^4 \frac{m_b m_\mu}{m_\sigma^2} \left[\frac{4x_{\pi_p}^2 - 7x_{\pi_p} + 1}{16(x_{\pi_p} - 1)^2} - \frac{x_{\pi_p}^2 - 2x_{\pi_p}}{8(x_{\pi_p} - 1)^3} \ln x_{\pi_p} \right], \\ C_{Q_2}(m_W)_{TC} &= -\frac{x_W}{\sin^2 \theta_W} \left(\frac{f}{f'} \right)^4 \frac{m_b m_\mu}{m_{\pi_p}^2} \left[\frac{x_{\pi_p} + 1}{8(x_{\pi_p} - 1)} - \frac{x_{\pi_p}}{4(x_{\pi_p} - 1)^2} \ln x_{\pi_p} \right]. \end{aligned} \quad (14)$$

From Eqs. (12-14) we find that (1) both the contributions arising from the neutral scalar exchange $C_{Q_{1,2}}$ and gauge boson exchange C_{10} are subject to helicity suppression, (2) the contributions arising from the neutral scalar exchanges are proportional to $(f/f')^4$, while those from the gauge bosons exchanges proportional to $(f/f')^2$. So for a sufficiently large f/f' , the contributions of neutral scalar exchanges are relatively enhanced and may become comparable with those from the gauge boson exchanges.

The Wilson coefficients at the lower scale of about m_b can be evaluated down from m_W scale by using the renormalization group equation. At leading order, the Wilson coefficients are [16, 17]

$$C_{10}(m_b) = C_{10}(m_W), \quad (15)$$

$$C_{Q_i}(m_b) = \eta^{-\gamma_Q/\beta_0} C_{Q_i}(m_W). \quad (16)$$

where $\beta_0 = 11 - 2n_f/3$, $\eta = \alpha_s(m_b)/\alpha_s(m_W)$ and $\gamma_Q = -4$ is the anomalous dimension of $\bar{q}P_R b$.

IV. CONSTRAINTS FROM R_b

Before presenting the numerical results, let us consider the current bounds on technicolor with scalars from a variety of experiments, especially the measurement of R_b . Using the Feynman rules in Ref. [8], one can easily find that the contributions from neutral scalars are negligible compared with those from charged scalars which appear in Fig. 1, and the bottom mass-dependent terms in R_b can also be omitted safely. In these approximations the addition contribution in the technicolor with scalars is obtained as

$$\delta R_b = R_b^{SM} (1 - R_b^{SM}) \Delta^{TC} \quad (17)$$




FIG. 1: Charged scalars diagrams contributing to $Zb\bar{b}$.

with

$$\begin{aligned} \Delta^{TC} &= \left(\frac{f}{f'} \right)^2 \frac{\alpha}{4\pi \sin^2 \theta_W} \frac{m_t^2}{m_W^2} \frac{v_{bL}}{v_{bL}^2 + v_{bR}^2} \{ v_{bL} B_1 \\ &\quad + v_{tR} \left[m_Z^2 (C_{22}^a - C_{23}^a) + 2C_{24}^a - \frac{1}{2} \right] \\ &\quad - 2v_{tL} m_t^2 C_0^a - \cos 2\theta_W C_{24}^b \}. \end{aligned} \quad (18)$$

Here $B_1 = B_1(-p_1, m_t, m_{\pi_p})$, $C_{0,ij}^a = C_{0,ij}(p_1, -P, m_{\pi_p}, m_t, m_t)$ and $C_{24}^b = C_{24}(-p_1, P, m_t, m_{\pi_p}, m_{\pi_p})$, with $p_1(p_2)$ and P denoting the four-momentum of $b(\bar{b})$ and Z boson respectively, are the Feynman loop integral functions and their expressions can be found in [18]. The coupling constants v_{qL} and v_{qR} are given by

$$v_{qL} = T_3^q - e_q \sin^2 \theta_W, \quad v_{qR} = -e_q \sin^2 \theta_W. \quad (19)$$

Our explicit expressions are not in agreement with those used in [6] where the results obtained in the framework of the two-Higgs doublet model (THMD) [19] were adopted directly. We checked the calculations and confirmed our results.

The current measurement of R_b reported by the LEP is $R_b^{expt} = 0.21646 \pm 0.00065$ [11]. Comparing with the SM value $R_b^{SM} = 0.21573 \pm 0.0002$, we obtained the constraints in h versus m_σ plane shown in Figs. 2 and 3. Although our explicit expression for R_b is different from that used in [20], a comparison of Fig. 2, Fig. 3 with Figure 1 in Ref. [20] suggests that there is not a qualitative change in the results plotted.

Our numerical results show that the constraint on f/f' from R_b is quite stringent, i.e., the ratio of f/f' must be smaller than 1.9 at 95% C. L., implying that the neutral scalars will not give dominate contributions to the processes of $B_{s,d} \rightarrow \mu^+ \mu^-$. Since previous studies did not comment on any restriction on f/f' arising from R_b , this is a new and interesting conclusion.

In Figs. 2 and 3 we also display the bounds from $B^0 - \bar{B}^0$ mixing and from the limits of Higgs masses [20]. In technicolor theories where the charged scalars couple to fermions in a similar pattern as in type-I two-Higgs doublet model, the strongest limit $m_{\pi_p^\pm} \geq 79$ GeV has been obtained directly from LEP experiments [12]. On the other hand, the LEP collaborations [12] have placed a 95% C. L. lower limit on the SM Higgs boson $M_H^0 \geq 113.5$




FIG. 2: Constraints on technicolor with scalars in limit (i). The allowed parameter space is the shaded region bounded by the contours $m_\sigma = 114$ GeV, δR_b (R_b line) and $hf' = 4\pi f$. The current bound from the searches for charged scalars $m_{\pi_p} = 79$ GeV is shown along with the reference curves $m_{\pi_p} = m_t - m_b$, $m_{\pi_p} = 1$ TeV. The constraint from $B^0 - \bar{B}^0$ mixing is labeled “B-line”.

GeV from searching for the process $e^+e^- \rightarrow Z^* \rightarrow ZH^0$. Although the limit on technicolor scalars may differ from that on M_H^0 , in practice, the contour $m_\sigma = 114$ GeV can serve as an approximate boundary to the experimentally allowed region [2, 20]. Note that the chiral Lagrangian analysis break down only constrain on the parameter space in limit (i) [6], the area above and to left of $hf' = 4\pi f$ line is excluded because the technifermion current masses are no longer small compared to the chiral symmetry breaking scale. For references, we also plotted the contours $m_{\pi_p} = m_t - m_b$ and $m_{\pi_p} = 1$ TeV. If the top quark doesn't decay to $\pi_p^+ b$, the areas *outside* of $m_{\pi_p} = m_t - m_b$ curve is excluded in Fig. 2. Similar situation occurs to $m_{\pi_p} = 1$ TeV curve in Fig. 3 if all scalar masses are restricted to the sub-TeV regime. In contrast to these, the excluded parameter space are the areas *inside* of $m_\pi = m_t - m_b$ curve in limit (ii) and $m_{\pi_p} = 1$ TeV curve in limit (i).

The constraint from $b \rightarrow s\gamma$ is close to that from $B^0 - \bar{B}^0$ mixing [7, 8, 21, 22], which are weaker than those from R_b [6]. As for the constraints from the measurement of $g_\mu - 2$, our previous study [23] showed that if the deviation of the E821 experiment result [14] and SM prediction $\Delta a_\mu \equiv a_\mu^{exp} - a_\mu^{SM} = (43 \pm 16) \cdot 10^{-10}$ persists, it would severely constrain the technicolor models because the technicolor models can hardly provide such a large contribution. However, over the last year the theoretical prediction of a_μ in the SM has undergone a




FIG. 3: Constraints on technicolor with scalars in limit (ii). The allowed parameter space is the shaded region bounded by the contours $m_\sigma = 114$ GeV and δR_b (R_b line). Other bound curves are the same as Fig. 2.

significant revision due to the change in sign of the light hadronic correction, which leads to only a 1.6σ deviation from the SM [24], yielding no more useful limits on this model.

V. NUMERICAL RESULTS

Bearing the constraints on technicolor with scalars in mind, and for the same values of m_{π_p} and f/f' , the allowed value of m_σ is generally smaller in limit (i), from Eq.(14) one can infer easily that the additional contributions to $B_{s,d} \rightarrow \mu^+\mu^-$ in limit (i) will be larger than those in limit (ii). Furthermore, as can be seen from the numerical coefficients in Eq. (12), the decay rate of B_s is significantly larger than B_d due primarily to the relative size of $|V_{ts}|$ to $|V_{td}|$. We thus take the B_s decay in limit (i) as an example to show the numerical results.

The experimental bound on $B_s \rightarrow \mu^+\mu^-$ comes from the CDF[25]

$$Br(B_s \rightarrow \mu^+\mu^-) < 2.6 \times 10^{-6} \quad (20)$$

at 95% C.L. with the corresponding integrated luminosity about $100 pb^{-1}$, while the SM prediction

$$Br(B_s \rightarrow \mu^+\mu^-) = 4.0 \times 10^{-9} \quad (21)$$

is obtained by taking the central values for all inputs in Eq. (12). The branching ratio of $B_s \rightarrow \mu^+\mu^-$ as a function of m_σ is displayed in Fig. 4 for various values of f/f' . The 2σ bounds at the upgraded Tevatron with $10 fb^{-1}$




FIG. 4: $Br(B_s \rightarrow \mu^+ \mu^-)$ as a function of m_σ for $f/f' = 1, 1.9, 5, 10$ (dash lines) in limit (i). The current 2σ upper bound [25] (dotted line), the SM prediction (solid line) as well as the expected sensitivity of the upgraded Tevatron with $10fb^{-1}$ and $20fb^{-1}$ (the dash-dotted lines) are also shown.

and $20fb^{-1}$ are also plotted under the assumption that the background for this decay is negligible. The corresponding expected sensitivity can be reach a branching

ratio of 1.3×10^{-8} and 6.5×10^{-9} (dash lines), respectively. We see that the R_b constraint $f/f' \leq 1.9$ at 95% C.L. shown in Fig. 4 is the strongest bound. Comparatively, the current upper bound on $Br(B_s \rightarrow \mu^+ \mu^-)$ from CDF [25] is much weaker, which only excludes a small region with large f/f' . Under the constraint $f/f' \leq 1.9$, the enhancement factor for the branching ratio in the technicolor model can still be up to 5. The upgraded Tevatron with $20fb^{-1}$ will be sensitive to enhancements of $B_s \rightarrow \mu^+ \mu^-$ in this model provided that m_σ is below 580 GeV.

VI. CONCLUSIONS

We have evaluated the decays $B_{s,d} \rightarrow \mu^+ \mu^-$ in the technicolor model with scalars, taking into account various experimental constraints, especially R_b , on the model parameter space. We first examined the restriction on f/f' arising from R_b which that previous study did not consider. We found that large f/f' , which might cause significantly enhancement for $Br(B_{s,d} \rightarrow \mu^+ \mu^-)$ from neutral scalars in the model, has been excluded by the constraints from R_b . Nevertheless, under the renewed R_b constraint, the branching ratio of $B_s \rightarrow \mu^+ \mu^-$ can still be enhanced by a factor of 5 relative to the SM prediction. With the maximum allowed value of $f/f' \sim 1.9$ from R_b , the upgraded Tevatron with $20fb^{-1}$ will be sensitive to enhancements of $B_s \rightarrow \mu^+ \mu^-$ in this model provided that m_σ is below 580 GeV. Since the theoretical uncertainties, which primarily come from the B-meson decay constants and CKM matrix elements, will be reduced in the on-going B-physics experiments and the lattice calculations, the processes $B_{s,d} \rightarrow \mu^+ \mu^-$ will promise to be a good probe of new physics.

-
- [1] E. H. Simmons, Nucl. Phys. B **312** (1989) 253.
 - [2] C.D. Carone and H. Georgi, Phys. Rev. D **49** (1994) 1427.
 - [3] R. S. Chivukula, S. B. Selipsky and E. H. Simmons, Phys. Rev. Lett. **69** (1992) 575.
 - [4] S. Samuel, Nucl. Phys. B **347** (1990) 625; M. Dine, A. Kagan, and S. Samuel, Phys. Lett. B **243** (1990) 250; A. Kagan and S. Samuel, Phys. Lett. B **252** (1990) 605.
 - [5] N. Evans, Phys. Lett. B **331** (1994) 378; C. D. Carone and E. H. Simmons, Nucl. Phys. B **397** (1993) 591; C.D. Carone and M. Golden, Phys. Rev. D **49** (1994) 6211.
 - [6] C. D. Carone, E. H. Simmons and Y. Su, Phys. Lett. B **344** (1995) 287.
 - [7] Y. Su, Phys. Rev. D **56** (1997) 335.
 - [8] Z. Xiong, H. Chen and L. Lu, Nucl. Phys. B **561** (1999) 3; Z. H. Xiong and J. M. Yang, Nucl. Phys. B **602** (2001) 289.
 - [9] For examples, see S. R. Choudhury and N. Gaur, Phys. Lett. B **451** (1999) 86; K. S. Babu and C. Koda, Phys. Rev. Lett. **84** (2000) 228; P. H. Chankowski and L. Slawianowska, Phys. Rev. D **63** (2001) 054012; C. Bobeth, T. Ewerth, F. Küger and J. Urban, Phys. Rev. D **64** (2001) 074014; G. D'Ambrosio, G. F. Giudice, G. Isidori and A. Strumia, hep-ph/0207036; C.-S. Huang, W. Liao, Q.-S. Yan and S.-H. Zhu, Phys. Rev. D **64** (2001) 05992; Z. H. Xiong and J. M. Yang, Nucl. Phys. B **628** (2002) 193; G. Isidori and A. Retico, JHEP **0111** (2001) 001; A. J. Buras, P. H. Chankowshi, J. Rosiek and L. Slawianowska, hep-ph/0207241. A. Dedes, H. K. Dreiner and U. Nierste, Phys. Rev. Lett. **87** (2001) 251804; A. Dedes, H. K. Dreiner, U. Nierste and P. Richardson, hep-ph/0207026.
 - [10] G. R. Lu *et al.*, Phys. Rev. D **54** (1996) 5647; Z. Phys. C **74** (1997) 355; G. R. Lu, Z. Xiong and Y. G. Cao, Nucl. Phys. B **487** (1997) 43; G. R. Lu, Z. Xiong, X. L. Wang and J. S. Huang, J. Phys. G **24** (1998) 745.
 - [11] T. Kawamoto, hep-ex/0105032; E. Tournefier, hep-ex/0105091; J. Drees, hep-ex/0110077.
 - [12] D.E. Groom *et al.*, Eur. Phys. J. C **15** (2000) 1.
 - [13] ALEPH Collaboration, Phys. Lett. B **429** (1998) 429; M. Nakao, Proceedings, ICHEP 2000, Osaka, Japan; T. Coan, Proceedings, ICHEP 2000, Osaka, Japan.
 - [14] H. N. Brown, *et al.*, Mu g-2 Collaboration,

- Phys. Rev. Lett.**86** (2001) 2227.
- [15] S. Coleman and E. Weinberg, Phys. Rev. **D7** (1973) 1888.
 - [16] B. Grinstein, R. Springer, and M. B. Wise, Phys. Lett. **B202** (1988) 138.
 - [17] Y. B. Dai, C. S. Huang and H. W. Huang, Phys. Lett. **B390** (1997) 257.
 - [18] G. 't Hooft and M. Veltman, Nucl. Phys. **B153** (1979) 365; G. Passarino and M. Veltman, Nucl. Phys. **B160** (1979) 151.
 - [19] M. Boulare and Donald Finnell, Phys. Rev. **D44** (1991) 2054 .
 - [20] V. Hemmige and E. H. Simmons, Phys. Lett. B **518** (2001) 72.
 - [21] B. Grinstein, R. Springer, and M. B. Wise, Nucl. Phys. **B339** (1990) 269.
 - [22] T. G. Rizzo, Phys. Rev. **D38** (1988) 820; W. S. Hou and R. S. Willey, Phys. Lett. **B202** (1988) 591 ; C. Q. Geng and J. N. Ng, Phys. Rev. **D38** (1988) 2858; V. Barger, J. L. Hewett, and R. Phillips, Phys. Rev. **D41** (1990) 3421.
 - [23] Z. H. Xiong and J. M. Yang, Phys. Lett. **B508** (2001) 295.
 - [24] M. Knecht and A. Nyffeler, Phys. Rev. **D65** (2002) 073034; M. Knecht, A. Nyffeler, M. Perrottet and E. de Rafael, Phys. Rev. Lett.**88** (2002) 071802; M. Hayakawa and T. Kinoshita, hep-ph/0112102; I. Blokland, A. Czarnecki and K. Melnikov, Phys. Rev. Lett.**88** (2002) 071803; J. Bijnens , E. Pallante and J. Prades, Nucl. Phys. **B626** (2002) 410.
 - [25] F. Abe *et. al.* (CDF collaboration), Phys. Rev. **D57** (1998) 3811.